



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2019

---

## **Current Aspects on Differentiating Relapses from Over-Infections in Symptomatic Inflammatory Bowel Diseases**

Liatsos, Christos ; Papaefthymiou, Apostolis ; Tzouvala, Maria ; Doulberis, Michael ; Petridou, Evangelia ; Kountouras, Jannis

DOI: <https://doi.org/10.1007/s10620-019-05755-8>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-181153>

Journal Article

Originally published at:

Liatsos, Christos; Papaefthymiou, Apostolis; Tzouvala, Maria; Doulberis, Michael; Petridou, Evangelia; Kountouras, Jannis (2019). Current Aspects on Differentiating Relapses from Over-Infections in Symptomatic Inflammatory Bowel Diseases. *Digestive Diseases and Sciences*, 64(9):2686-2687.

DOI: <https://doi.org/10.1007/s10620-019-05755-8>

# Current Aspects on Differentiating Relapses from Over-Infections in Symptomatic Inflammatory Bowel Diseases

Christos Liatsos<sup>1,6\*</sup> Apostolis Papaefthymiou<sup>1</sup> Maria Tzouvala<sup>2</sup> Michael Doulberis<sup>3,5</sup> Evangelia Petridou<sup>4</sup> Jannis Kountouras<sup>5</sup>

1 Department of Gastroenterology, 401 General Military Hospital of Athens, 11525 Athens, Attiki, Greece

2 Department of Gastroenterology, Agios Panteleimon General Hospital, 18454 Nikaia, Attiki, Greece

3 Department of Gastroenterology and Hepatology, University of Zurich, 8091 Zurich, Switzerland

4 Department of Microbiology, “Agia Sofia” Pediatric Hospital, 11527 Athens, Attiki, Greece

5 Department of Internal Medicine, Second Medical Clinic, Ippokration Hospital, Aristotle University of Thessaloniki, 54642 Thessaloniki, Macedonia, Greece

6 PO Box 833, 19009 Pikermi, Rafina, Greece

\* Christos Liatsos [cliatsos@yahoo.com](mailto:cliatsos@yahoo.com)

Dear Editor,

In their comparative retrospective cohort, Ahmad et al. [1] showed a higher rate of adverse events such as hospitalization, surgery, or delayed inflammatory bowel disease (IBD) treatment modification in patients with IBD tested positive with stool multiplex gastrointestinal pathogen panels (GPPs) for discrimination of diarrhea related to IBD exacerbation or infection, though positive predictive value of GPPs remained limited [1]. Gut microbiota of IBD patients includes a plethora of infectious species that could cause false positive results in GPP tests, thus delaying the decision of the appropriate treatment and reducing its clinical benefit [2]. Moreover, the interpretation of GPP test results in clinical practice is associated with some limitations, also mentioned by authors [1]. In this respect, several pathogens can occur not only asymptotically (e.g., norovirus and *Salmonella* spp.) but also subclinically (e.g., *Clostridium difficile* non-toxigenic strains) in a colonization-like setting [3, 4], where connection with disease is uncertain. Additionally, an uncertainty exists regarding the cost-effectiveness of GPPs for suspected infectious gastroenteritis in hospital and community settings [5]. Interestingly, despite the well-described biases of GPPs in IBD [1] especially in terms of antimicrobial stewardship [6], all of the panels lack the capability to detect cytomegalovirus (CMV), which consists a substantial burden of IBD flares and steroids refractoriness being detected in 21–34% and 33–36% of these cases, respectively

[7]. Nevertheless, even if GPPs were able to detect fecal CMV-DNA, the sensitivity of polymerase chain reaction (PCR) for CMV-genome in stools seems to remain inferior to its detection in tissue samples depending also on the viral load [8]. Another important aspect that should be introduced is the role of confocal endomicroscopy (CE), which rises in IBD flare identification, as specific characteristics to define IBD activity have been suggested in many studies. Although still experimental, the assessment of the crypt morphology (number of colonic crypts, crypt tortuosity, crypt lumen), erosions, vascularity, cellular infiltrates within the lamina propria, number of goblet cells, gut barrier disruption, and leakage of fluorescein to the lumen could detect immediately and non-invasively the etiology of symptom exacerbation among IBD patients [9–11]. Furthermore, *Clostridium difficile* is visible and recognizable in vivo in the gut mucosa under CE, irrespective of symptoms manifestation [12], whereas other over-infections in patients with IBD have not been evaluated using CE, although it is known that these patients have intramucosal presence of bacteria more frequently than normal controls [9]. Nonetheless, this technique is still limited, lacking evidence-based data, expensive, and difficult to be performed to replace the conventional laboratory workup. Recently, by using the Luminex xTAG GPP, quantitative thresholds offer recognition of *Clostridium difficile* infection [13] which has been increasingly associated with IBD flares [14]. As deduced by the clinical necessity to differentiate IBD flares from over-infections beyond multiplex GPPs, further studies are required to develop and assess a novel, validated, and easy-to-perform, grading diagnostic index including clinical, laboratory, microbiological, and endoscopic findings to increase the sensitivity of a secure and accurate diagnosis in IBD symptomatic patients and obtain the potentiality of an immediate intervention.

## **Compliance with Ethical Standards**

Conflict of interest: The authors declared that they have no conflict of interest.

## **References**

1. Ahmad W, Nguyen NH, Boland BS, et al. Comparison of multiplex gastrointestinal pathogen panel and conventional stool testing for evaluation of diarrhea in patients with inflammatory bowel diseases. *Dig Dis Sci*. 2019;64:382–390. <https://doi.org/10.1007/s10620-018-5330-y>

2. Ni J, Wu GD, Albenberg L, Tomov VT. Gut microbiota and IBD: causation or correlation? HHS public access. *Nat Rev Gastroenterol Hepatol*. 2017;14:573–584.
3. Eyre DW, Griffiths D, Vaughan A, et al. Asymptomatic *clostridium difficile* colonisation and onward transmission. *PLoS ONE*. 2013;8:e78445.
4. Teunis PFM, Sukhrie FHA, Vennema H, Bogerman J, Beersma MFC, Koopmans MPG. Shedding of norovirus in symptomatic and asymptomatic infections. *Epidemiol Infect*. 2015;143:1710–1717.
5. Freeman K, Mistry H, Tsertsvadze A, et al. Multiplex tests to identify gastrointestinal bacteria, viruses and parasites in people with suspected infectious gastroenteritis: a systematic review and economic analysis. *Health Technol Assess (Rockv)*. 2017;21:1–188.
6. Fishman N. Antimicrobial stewardship. *Am J Med*. 2006;119:S53– S61. (discussion S62-70).
7. Johnson J, Affolter K, Boynton K, Chen X, Valentine J, Peterson K. CMV Disease in IBD: comparison of diagnostic tests and correlation with disease outcome. *Inflamm Bowel Dis*. 2018;24:1539–1546.
8. Thörn M, Rorsman F, Rönnblom A, et al. Active cytomegalovirus infection diagnosed by real-time PCR in patients with inflammatory bowel disease: a prospective, controlled observational study. *Scand J Gastroenterol*. 2016;51:1075–1080.
9. Neumann H, Vieth M, Atreya R, et al. Assessment of Crohn's disease activity by confocal laser endomicroscopy. *Inflamm Bowel Dis*. 2012;18:2261–2269.
10. Kiesslich R, Duckworth CA, Moussata D, et al. Local barrier dysfunction identified by confocal laser endomicroscopy predicts relapse in inflammatory bowel disease. *Gut*. 2012;61:1146–1153.
11. Rasmussen DN, Karstensen JG, Riis LB, Brynskov J, Vilmann P. Confocal laser endomicroscopy in inflammatory bowel disease—a systematic review. *J Crohn's Colitis*. 2015;9:1152–1159.
12. Neumann H, Günther C, Vieth M, et al. Confocal laser endomicroscopy for in vivo diagnosis of *clostridium difficile* associated colitis—a pilot study. *PLoS ONE*. 2013;8:1–7.
13. Leal SM, Popowitch EB, Levinson KJ, et al. Quantitative thresholds enable accurate identification of *clostridium difficile* infection by the Luminex xTAG gastrointestinal pathogen panel. *J Clin Microbiol*. 2018;56:e01885-17.

14. Khanna S, Shin A, Kelly CP. Management of clostridium difficile infection in inflammatory bowel disease: expert review from the clinical practice updates committee of the AGA institute. Clin Gastroenterol Hepatol. 2017;15:166–174.